

(19) World Intellectual Property  
Organization  
International Bureau



(43) International Publication Date  
29 July 2004 (29.07.2004)

PCT

(10) International Publication Number  
WO 2004/063146 A1

(51) International Patent Classification<sup>7</sup>: C07C 259/10,  
A61K 31/16, A61P 35/00, 29/00

(21) International Application Number:  
PCT/IT2004/000002

(22) International Filing Date: 7 January 2004 (07.01.2004)

(25) Filing Language: Italian

(26) Publication Language: English

(30) Priority Data:  
MI2003A000025 10 January 2003 (10.01.2003) IT

(71) Applicant (for all designated States except US): ITAL-  
FARMACO SPA [IT/IT]; Viale Fulvio Testi, 330, I-20126  
Milan (IT).

(72) Inventors; and

(75) Inventors/Applicants (for US only): PINORI, Massimo  
[IT/IT]; Via Edison, 24, I-23877 Paderno d'Adda (IT).  
MASCAGNI, Paolo [IT/IT]; Via don Galli, 25, I-20058  
Villasanta (IT). MAZZAFERRO, Rocco [IT/IT]; Via  
Pietro Pomponazzi, 19/B, I-20141 Milano (IT). VER-  
GANI, Barbara [IT/IT]; Via Laghetto, 24, I-20050  
Macherio (IT).

(74) Agents: PISTOLESI, Roberto et al.; Dragotti & Asso-  
ciati SRL, Galleria San Babila, 4/C, I-20122 Milano (IT).

(81) Designated States (unless otherwise indicated, for every  
kind of national protection available): AE, AG, AL, AM,  
AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN,  
CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI,  
GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE,  
KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD,  
MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG,  
PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM,  
TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM,  
ZW.

(84) Designated States (unless otherwise indicated, for every  
kind of regional protection available): ARIPO (BW, GH,  
GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),  
Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), Euro-  
pean (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR,  
GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK,  
TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,  
ML, MR, NE, SN, TD, TG).

**Declaration under Rule 4.17:**


— of inventorship (Rule 4.17(iv)) for US only

**Published:**

— with international search report  
— before the expiration of the time limit for amending the  
claims and to be republished in the event of receipt of  
amendments

For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.

(54) Title: HYDROXAMIC ACID DERIVATIVES HAVING ANTI-INFLAMMATORY ACTION



(57) Abstract: Compounds of formula (I) are described: The compounds (I) inhibit TNF $\alpha$  production and may therefore be useful in the treatment of inflammation, of auto-immune diseases, and of pathological conditions which involve excessive production of that cytokine. The compounds (I) are also inhibitors of the histone deacetylase enzyme and may therefore be useful in tumorous diseases, alone or in association with other anti-tumour active ingredients.

**Title:**

Hydroxamic acid derivatives having anti-inflammatory action

**Description****Field of the invention**

The present invention relates to derivatives of hydroxamic acid, in particular derivatives of N-hydroxy benzamide, with anti-inflammatory and anti-tumour activity.

**Background of the invention**

Derivatives of hydroxamic acid with anti-inflammatory and immunosuppressive activity are described in US patent 6,034,096. These compounds can inhibit the production of pro-inflammatory cytokines, in particular of TNF $\alpha$  (tumour necrosis factor) and of interleukin-1-beta and can therefore be used in the treatment of conditions which involve excessive production of those substances, such as inflammatory and auto-immune diseases or tumorous forms.

The compounds described in US patent 6,034,096 are characterized in that they contain two cyclic structures which are linked by a carbamate or urea group and one of which is in turn linked to an N,hydroxy-carboxamide (hydroxamic acid) group.

**Summary of the invention**


It has now been found that, by introducing suitable substituent groups, other than those described in US 6,034,096, into the first cyclic structure, it is possible to modulate the potency of the TNF $\alpha$  inhibitory activity and to obtain compounds which are metabolically stable and only slightly cytotoxic. These compounds can therefore usefully be used for the treatment of inflammatory and/or auto-immune diseases

in which the production of TNF $\alpha$  and of other pro-inflammatory cytokines performs a pathological role.

It has also been found that the compounds of the present invention are inhibitors of histone deacetylase (HDAC) enzymes and, as such, can also usefully be used in the treatment of various tumorous, auto-immune or neurodegenerative pathological conditions, for which the inhibition of those enzymes is useful.

#### Description of the invention

The subject of the present invention is derivatives of hydroxamic acid of formula (I):



(I)

in which

R is hydrogen, C<sub>1-4</sub> alkyl or phenyl;

R' is hydrogen or C<sub>1-4</sub> alkyl;

A is aryl or a monocyclic, dicyclic, or tricyclic residue, optionally partly or completely unsaturated, containing one or more heteroatoms selected from the group formed by N, S and O and possibly substituted with haloalkyl, alkylsulphonyl, (cycloalkyl)alkyl, alkanoyl, carbonyloxy, alkoxy, thioalkoxy, thiophenoxy, nitro, cyano, oxo, perfluoroalkoxy, perfluoroalkyl, phenyl, phenoxy, phenylalkoxy, benzoyloxy, phenylalkyl, benzoyl, phenylsulphonyl groups, the phenyl, phenoxy, phenylalkoxy, phenylalkyl, benzoyl and benzoyloxy substituents optionally being substituted in the aromatic ring with alkyl, alkoxy, alkylsulphonyl, amino, cyano, hydroxy, nitro, perfluoroalkoxy,

perfluoroalkyl, phenylsulphonyl, thioalkoxy and halogen groups;

L is a chain of from 1 to 5 carbon atoms optionally containing a double bond or an NR' group in which R' is as defined above;

X is an oxygen atom, an NR' group in which R' is as defined above, or is absent;

r and m are, independently, 0, 1 or 2;

B is phenyl or cyclohexyl.

Preferred compounds according to the present invention are those in which:

R is hydrogen or methyl, more preferably hydrogen,

A is phenyl, preferably substituted with one or more groups selected from: alkoxy, nitro, perfluoroalkyl, phenoxy, phenyl, phenylalkoxy, benzoyloxy, thioalkoxy;

L is methylene or ethylene and X is absent;

m and r are equal to zero;

B is phenyl;

R' is hydrogen.

Moreover, the following compounds are particularly preferred:

N-hydroxy-4-[2-(4-trifluoromethyl-phenyl)-acetyl-amino]-benzamide;

N-hydroxy-4-(3-phenyl-butyrylamino)-benzamide;

N-hydroxy-4-[3-(3-methoxy-phenyl)-propionyl-amino]-benzamide;

N-hydroxy-4-[2-(4-methoxy-phenyl)-acetyl-amino]-benzamide;

N-hydroxy-4-[2-(4-ethoxy-phenyl)-acetyl-amino]-benzamide;

N-hydroxy-4-[3-(3,5-bis-trifluoromethyl-phenyl)-propionyl-amino]-benzamide;

N-hydroxy-4-[2-(3,4,5-trimethoxy-phenyl)-acetyl-amino]-

benzamide;  
N-hydroxy-4-[2-(4-methylsulphanil-phenyl)-acetylamino]-  
benzamide;  
N-hydroxy-4-[2-(3-trifluoromethyl-phenyl)-acetylamino]-  
benzamide;  
N-hydroxy-4-[2-(3-nitro-phenyl)-acetylamino]-benzamide;  
N-hydroxy-4-[2-(3-phenoxy-phenyl)-acetylamino]-  
benzamide;  
N-hydroxy-4-[2-(diphenyl-4-yl)-acetylamino]-benzamide;  
N-hydroxy-4-[2-(2,3-dimethoxy-phenyl)-acetylamino]-  
benzamide;  
2-[2-(4-hydroxycarbamoyl-phenylcarbamoyl)-ethyl]-phenyl  
ester of benzoic acid;  
N-hydroxy-4-[2-(4-nitro-phenyl)-acetylamino]-benzamide;  
N-hydroxy-4-[2-(2-phenoxy-phenyl)-acetylamino]-  
benzamide;  
N-hydroxy-4-[2-(4,5-dimethoxy-2-nitro-phenyl)-  
acetylamino]-  
benzamide;  
N-hydroxy-4-[2-(2-benzyloxy-phenyl)-acetylamino]-  
benzamide;

N-hydroxy-4-[2-(2-nitro-phenyl)-acetylamino]-benzamide.  
Compounds of general formula (I), and this is a first  
aspect of the invention, can inhibit *in vitro* and *in vivo*,  
at nanomolar concentrations, the production of  
TNF $\alpha$  and of other pro-inflammatory cytokines.

A second aspect of the present invention consists of  
the use of compounds of the general formula (I), alone  
or in combination with other active ingredients, for  
the treatment of inflammatory and auto-immune  
pathological conditions which are characterized by  
excessive production of TNF $\alpha$  and/or of other pro-  
inflammatory cytokines such as, for example,

spondyloarthropathy (*Expert Opin. Emerging Drugs* (2002) 7(2):235-246), rheumatoid arthritis (*Lancet* (1999) 354(9194): 1932), acute alcoholic hepatitis (*Am. J. Gastroenterology* (2001) 96(12):3361-3367), inflammatory syndromes of the intestine such as Crohn's disease and ulcerative colitis (*Drugs of Today* (2000) 36(5):281-293), heart failure (*Heart Fail. Rev.* (2001) 6(2): 143), asthma (*Int. J. Biochem. Cell Biol.* (2000) 32(8): 833), intracerebral haemorrhage (*Stroke* (2001) 32:240-248), psoriasis (*Drugs Today* (1999) 35(12):913), diabetes (*J. Autoimmun.* (2003) 20(4):303-312 - *Pancreas*. (2003) 26(4):E99-E104), etc.

In a third aspect of the invention, the compounds of general formula (I) are inhibitors of the activity of the histone deacetylase enzymes.

A further aspect of the invention therefore consists of the use of compounds of general formula (I), alone or in combination with other active ingredients, for the treatment of tumorous pathological conditions (*TRENDS in Endocrinology & Metabolism* (2001) 12 (7):294-300), neurodegenerative pathological conditions (*Nature* (2001) 413:739-743), and auto-immune pathological conditions (*J. Clinical Investigation* (2003) 111:539-552).

The compounds (I) are also inhibitors of the histone deacetylase enzyme and may therefore be useful in tumorous diseases, alone or in combination with other anti-tumour active ingredients; to maximize the anti-tumour effect, the compounds (I) are preferably administered in combination with anti-tumour active ingredients having an action mechanism other than HDAC


inhibition, such as, for example, anti-proliferative active ingredients.

Pharmaceutical compositions which contain compounds of formula (I) are therefore also subjects of the present invention; these compositions may be in the form of capsules, tablets, coated tablets, creams, ointments or phials for oral, intramuscular, or intravenous administration.

In the above-mentioned pharmaceutical compositions, the compounds of formula (I), alone or in combination with other active ingredients, may optionally be mixed with conventional excipients or vehicles, for example, those described in Remington's Pharmaceutical Sciences Handbook, XVII, ed. Mack Pub., N.Y., U.S.A.


The compounds of formula (I) may be prepared by the method described in US 6,034,096, which is incorporated herein by reference. Alternatively, the compounds of the invention may be prepared by "solid phase" organic synthesis with the use of one of the commercially available special resins for N-hydroxyamides. For example, a polystyrene resin cross-linked with divinyl benzene, functionalized with O-alkylated hydroxylamine groups from para-alkoxybenzyl residues (Wang type resins) may be used [see e.g. Richter, L.S. and Desai M.C., *Tetrahedron Letters* (1996) 38(3): 321-322].

The amine groups that are present in the resin may be acylated by suitably protected amino-acids in the presence of suitable condensing agents to give intermediates of formula



where G is a suitable protector group and m, r and B have the meanings defined for the general formula (I).

After removal of the protector group G, the amine group can be further acylated with the use, after activation or in the presence of suitable condensing agents, of acids of formula (III):



where R, A, L and X have the meanings defined for the general formula (I).

Finally, the products of the invention can be released from the resin by treatment with medium strength acids (for example, trifluoroacetic acid), filtration and possibly final purification.

The present invention will be illustrated below by means of some examples which have the purpose purely of further describing the subjects of the present invention and are not intended as in any way limiting thereof.

#### EXAMPLES

The following abbreviations are used in the examples given below:

ACN	acetonitrile
PVDF	polyvinylidene difluoride
DCM	dichloromethane
DMF	dimethyl formamide
HOAt	1-hydroxy-7-azabenzotriazole
HATU	O-(7-azabenzotriazol-1-yl)-N,N,N'-tetramethyl-
	uronium hexafluorophosphate
TFA	trifluoroacetic acid
General purification method	



Unless indicated otherwise, all of the final purifications were performed by a Waters HPLC/MS preparation system with a Waters Symmetry C18 5  $\mu$ m 19x50 mm column equipped with a Waters ZQ mass spectrometer.

Operative conditions:

ES+ centroid ionization, scanning time 15 min., scanning m/z 120-1000, cone voltage 15V, source temperature 120°C, solvation temperature 250°C.

HPLC eluents:

A=H<sub>2</sub>O, B=ACN, C=HCOOH 1% in H<sub>2</sub>O

Gradient:

Time (min)	A	B	C	Flow (ml/min)
0	94%	5%	1%	20
2	94%	5%	1%	20
3	87%	12%	1%	20
8	87%	12%	1%	20
11	20%	80%	1%	20
12	94%	5%	1%	20

An aliquot of the crude product to be purified (30-50 mg) was dissolved in 0.1 ml of MeOH and diluted with 0.4 ml of ACN/H<sub>2</sub>O mixture (1:1; v/v). The solution was filtered on 0.45 µm PVDF membrane and was injected into the preparation system described above. For each run, the fractions corresponding to the peak associated with the expected molecular ion ([M+H]<sup>+</sup>) were collected, recombined and concentrated to dryness.

#### Example 1

**N-hydroxy-4-[2-(4-trifluoromethyl-phenyl)-acetylaminol]-benzamide**

##### *Step A*

A mixture of HOAt (55 mg, 0.4 mmoles) and HATU (152 mg, 0.4 mmoles) in anhydrous DMF (0.5 ml), followed by diisopropyl ethylamine (0.14 ml, 0.8 mmoles), was added to a solution of 4-(9H-fluoren-9-yl-methoxycarbonylamino)-

benzoic acid (150 mg, 0.4 mmoles) in anhydrous DMF (0.5 ml). The reaction mixture was stirred at ambient temperature for 30 minutes and was then transferred into a reactor containing a Wang type polystyrene resin, functionalized with hydroxylamine (0.1 g; 0.1 mmoles), and the suspension was stirred at ambient temperature for 16 hours. The resin was filtered and washed, in sequence, with DMF (5x2 ml), DCM (2x2 ml), MeOH (2x2 ml), DCM (2x2 ml), MeOH (2x2 ml), DCM (2x2 ml) and DMF (5x2 ml); finally, the resin was filtered and dried under vacuum.

#### Step B

The resin obtained in A was expanded in a 20% solution of piperidine in DMF (1 ml) and stirred at ambient temperature for one hour and was then filtered, washed with DMF (5x2 ml), and dried under vacuum.

#### Step C

HATU (381 mg, 1 mmole) and diisopropyl ethylamine (262  $\mu$ l, 1.5 mmoles) were added to a solution of (4-trifluoromethyl-phenyl)-acetic acid (184 mg, 0.9 mmoles) in anhydrous DMF (1 ml). The reaction mixture was stirred at ambient temperature for 30 minutes and was then added to the resin obtained in B and stirred again at ambient temperature for 20 hours. The resin was filtered and washed, in sequence, with DMF (5x2 ml), DCM (2x2 ml), MeOH (2x2 ml), DCM (2x2 ml), MeOH (2x2 ml), DCM (5x2 ml) and, finally, was filtered and dried under vacuum.

#### Step D

The resin obtained in C was expanded in a 50% solution of TFA in DCM (1 ml) and was subjected to stirring at ambient temperature for one hour and was then filtered and the solution was evaporated to dryness. The

residue was taken up with t-BuOMe and re-evaporated to dryness five times. The crude product thus obtained was purified by preparation HPLC/MS, following the general method described above.

10.5 mg was obtained;  $[M+H]^+ = 339.3$  (calc. 339.1)

#### Example 2

##### **N-hydroxy-4-(3-phenyl-butyrylamino)-benzamide**

The product was prepared by the method described in Example 1, with the use of 3-phenyl-butyric acid in step C.

8.9 mg was obtained;  $[M+H]^+ = 299.3$  (calc. 299.1)

#### Example 3

##### **N-hydroxy-4-[3-(3-methoxy-phenyl)-propionylamino]-benzamide**

The product was prepared by the method described for Example 1, with the use of 3-(3-methoxy-phenyl)-propionic acid in step C.

15.7 mg was obtained;  $[M+H]^+ = 315.3$  (calc. 315.1)

#### Example 4

##### **N-hydroxy-4-[2-(4-methoxy-phenyl)-acetylaminol]-benzamide**

##### **Step A**

4-methoxy-phenyl acetic acid (5g) was suspended in chloroform (150 ml); thionyl chloride (10.8 g) was added to the suspension and was heated to reflux, giving a clear solution. After four hours under reflux, the solvent was evaporated under vacuum and the residue was taken up with further chloroform and evaporated again. The residue was dissolved in tetrahydrofuran (100 ml) and p-amino-benzoic acid (4.1 g) dissolved in tetrahydrofuran (100 ml) was added to the solution. The reaction mixture was left at ambient temperature overnight. The reaction mixture was then

evaporated under vacuum and the residue was crystallized from 70% ethanol (250 ml). 5.1 g of product was obtained (yield 60%) with HPLC purity greater than 97%.

$[M+H]^+ = 286.3$  (calc. 286.1)

#### Step B

The product obtained in the preceding step (5 g) was suspended in chloroform (150 ml); thionyl chloride (6.3 g) was added to the suspension and was heated to reflux giving a clear solution. After four hours under reflux, the solvent was evaporated under vacuum and the residue was taken up with further chloroform and evaporated again. The residue was dissolved in tetrahydrofuran (100 ml) and added to an aqueous hydroxylamine solution (50% w/w; 6.5 ml) in tetrahydrofuran (100 ml). The reaction mixture was left at ambient temperature overnight. The reaction mixture was then evaporated under vacuum and the residue was crystallized from 70% ethanol (100 ml). 4.14 g of product was obtained (yield 78%) with HPLC purity greater than 99%; m.p.=197.6-200°C (dec.);  $[M+H]^+ = 301.3$  (calc. 301.1)

#### Example 5

##### **N-hydroxy-4-[2-(4-ethoxy-phenyl)-acetylamino]-benzamide**

The product was prepared by the method described for Example 1, with the use of 2-(4-ethoxy-phenyl)-acetic acid in step C.

14.9 mg was obtained;  $[M+H]^+ = 315.3$  (calc. 315.1)

#### Example 6

##### **N-hydroxy-4-[3-(3,5-bis-trifluoromethyl-phenyl)-propionyl-amino]-benzamide**

The product was prepared by the method described for Example 1, with the use of 3-(3,5-bis-trifluoromethyl-phenyl)-propionic acid in step C.

14.9 mg was obtained;  $[M+H]^+=421.3$  (calc. 421.1)

Example 7

**N-hydroxy-4-[2-(3,4,5-trimethoxy-phenyl)-acetylamino]-benzamide**

The product was prepared by the method described for Example 1, with the use of 2-(3,4,5-trimethoxy-phenyl)-acetic acid in step C.

11.4 mg was obtained;  $[M+H]^+=361.4$  (calc. 361.1).

Example 8

**N-hydroxy-4-[2-(4-methylsulphanil-phenyl)-acetylamino]-benzamide**

The product was prepared by the method described for Example 1, with the use of 2-(4-methylsulphanil-phenyl)-

acetic acid in step C.

10.6 mg was obtained;  $[M+H]^+=317.4$  (calc. 317.1)

Example 9

**N-hydroxy-4-[2-(3-trifluoromethyl-phenyl)-acetylamino]-benzamide**

The product was prepared by the method described for Example 1, with the use of 2-(3-trifluoromethyl-phenyl)-acetic acid in step C.

7.9 mg was obtained;  $[M+H]^+=339.3$  (calc. 339.1)

Example 10

**N-hydroxy-4-[2-(3-nitro-phenyl)-acetylamino]-benzamide**

The product was prepared by the method described for Example 1, with the use of 2-(3-nitro-phenyl)-acetic acid in step C.

16.1 mg was obtained;  $[M+H]^+=316.3$  (calc. 316.1)

Example 11

**N-hydroxy-4-[2-(3-phenoxy-phenyl)-acetylamino]-benzamide**

The product was prepared by the method described for Example 1, with the use of 2-(3-phenoxy-phenyl)-acetic acid in step C.

14.3 mg was obtained;  $[M+H]^+=363.4$  (calc. 363.1)

Example 12

**N-hydroxy-4-[2-(diphenyl-4-yl)-acetylamino]-benzamide**

The product was prepared by the method described for Example 1, with the use of 2-(diphenyl-4-yl)-acetic acid in step C.

9.1 mg was obtained;  $[M+H]^+=347.3$  (calc. 347.1)

Example 13

**N-hydroxy-4-[2-(2,3-dimethoxy-phenyl)-acetylamino]-benzamide**

The product was prepared by the method described for Example 1, with the use of 2-(2,3-dimethoxy-phenyl)-acetic acid in step C.

10.8 mg was obtained;  $[M+H]^+=331.3$  (calc. 331.1)

Example 14

**2-[2-(4-hydroxycarbamoyl-phenylcarbamoyl)-ethyl]-phenyl ester of benzoic acid**

The product was prepared by the method described for Example 1, with the use of 2-(2-carboxy-ethyl)-phenyl ester of benzoic acid in step C.

18.1 mg was obtained;  $[M+H]^+=405.4$  (calc. 405.1)

Example 15

**N-hydroxy-4-[2-(4-nitro-phenyl)-acetylamino]-benzamide**

The product was prepared by the method described for Example 1, with the use of 2-(4-nitro-phenyl)-acetic acid in step C.

7.2 mg was obtained;  $[M+H]^+=316.3$  (calc. 316.1)

Example 16

**N-hydroxy-4-[2-(2-phenoxy-phenyl)-acetyl-amino]-benzamide**

The product was prepared by the method described for Example 1, with the use of 2-(2-phenoxy-phenyl)-acetic acid in step C.

14.2 mg was obtained;  $[M+H]^+=363.4$  (calc. 363.1)

Example 17

**N-hydroxy-4-[2-(4,5-dimethoxy-2-nitro-phenyl)-acetyl-amino]-benzamide**

The product was prepared by the method described for Example 1, with the use of 2-(4,5-dimethoxy-2-nitro-phenyl)-acetic acid in step C.

8.6 mg was obtained;  $[M+H]^+=376.3$  (calc. 376.1)

Example 18

**N-hydroxy-4-[2-(2-benzyloxy-phenyl)-acetyl-amino]-benzamide**

The product was prepared by the method described for Example 1, with the use of 2-(2-benzyloxy-phenyl)-acetic acid in step C.

10.1 mg was obtained;  $[M+H]^+=377.4$  (calc. 377.1)

Example 19

**N-hydroxy-4-[2-(2-nitro-phenyl)-acetyl-amino]-benzamide**

The product was prepared by the method described for Example 1, with the use of 2-(2-nitro-phenyl)-acetic acid in step C.

12.5 mg was obtained;  $[M+H]^+=316.3$  (calc. 316.1)

Example 20

**Inhibition of TNF $\alpha$  production - determination *in vitro***

The compounds were dissolved in DMSO to a final concentration of 120 mM and were stored at -80°C. The solutions for the test were prepared by diluting the



mother solutions in RPMI 1640 with the addition of 1% FCS and 0.01% DMSO and were filtered with 0.2  $\mu$ m filters.

Mononuclear peripheral blood cells were obtained from "buffy coats" of healthy donors by separation in a Ficoll-Hypaque gradient. The cells were spread in plates with 96 wells at a concentration of about 500,000 cells per well, suspended in RPMI 1640 containing 1% FCS and were incubated at 37°C in the presence of various concentrations (from  $10^{-6}$  to  $10^{-11}$  M) of the compounds to be tested.

After 1 hour, LPS was added (to a final concentration of 10 ng/ml; obtained from *E. coli* 055:B5) and the plates were incubated at 37°C for a further 24 hours. Upon completion, the supernatant fluids were collected and used for the determination of the TNF $\alpha$  content by ELISA (ELISA DuoSet Kit; R&D systems, Minneapolis, MN, USA).

The concentration of TNF $\alpha$  was calculated with the use of a calibration curve and the IC<sub>50</sub> values (concentration which inhibits the production of the cytokine by 50%) were calculated from the curve obtained, giving the percentage inhibition values for each individual concentration of the compound under test.

The values obtained for the compounds described in the preceding examples are given in the following table (the values are the average of the results obtained in at least two determinations, performed with cells of different donors).

Table 1: Inhibition of TNF $\alpha$  production in human monocytes stimulated with LPS

Example	IC <sub>50</sub> (nM)	Example	IC <sub>50</sub> (nM)
Ex. 01	14.5	Ex. 11	108.0
Ex. 02	76.0	Ex. 12	227.0
Ex. 03	26.0	Ex. 13	187.0
Ex. 04	51.5	Ex. 14	157.0
Ex. 05	32.0	Ex. 15	27.0
Ex. 06	17.0	Ex. 16	259.0
Ex. 07	45.0	Ex. 17	280.0
Ex. 08	320.0	Ex. 18	391.0
Ex. 09	70.0	Ex. 19	1000.0
Ex. 10	134.0		

Example 21

**Inhibition of TNF $\alpha$  production - determination *in vivo***  
For the compounds of the present invention, the capacity to inhibit TNF $\alpha$  production induced by the administration of LPS was evaluated in mice.

A lethal quantity of LPS (*E. coli* 055:B5; 2 mg/kg) was administered to the animals (CD1 female mice) by an intraperitoneal route; 90 min. after administration, the animals were killed and the TNF $\alpha$  content present in the blood was determined by ELISA assay.

The compounds under test, suspended in Methyl Cellosolve (0.5% in water), were administered orally, 60 min. prior to the administration of LPS, at a dose of 1 mg/kg.

The results are given in the following table; the values are expressed as percentage inhibition of TNF $\alpha$  production in comparison with the control group.

Table 2: inhibition of TNF $\alpha$  production in mice

Example	Inhibition
Ex. 01	67 %
Ex. 03	56 %
Ex. 04	42 %

Example 22**Metabolic resistance *in vitro***

The metabolic resistance of some compounds described in the preceding examples was evaluated by incubating the substances with the S9 fraction of a microsomal preparation of liver cells. Each compound (6 µg/ml) was incubated at 37°C for 30 min. in phosphate buffer (pH 7.4) containing the S9 fraction (protein content 2 mg/ml). The reaction was stopped by cooling in an ice bath and adding an equal volume of water/acetonitrile (50:50) containing 0.2% of trifluoroacetic acid. After centrifuging, an aliquot of the supernatant fluid was analyzed by HPLC to evaluate the percentage of product that had remained intact.

The percentages of unchanged product that were present after incubation for 30 min. are given in the following table.

Table 3: metabolic transformation by means of the S9 liver fraction

Examples	Residual product after 30 min.
Ex. 01	78.9 %
Ex. 03	60.7 %
Ex. 04	90.2 %
Ex. 05	89.0 %

Ex. 06	68.4 %
--------	--------

Example 23**Cytotoxicity *in vitro***

The cytotoxicity of some compounds described in the preceding examples was evaluated *in vitro* on the human HEP-G2 hepatoma cell line by a commercial colorimetric method (Cell Titer 96® Aqueous One Solution Cell Proliferation Assay - Promega); the method determines the number of living cells on the basis of their ability to metabolize a tetrazolium salt producing formazane. The quantity of formazane produced is proportional to the number of living cells.

The HPE-G2 cells were distributed in micro-plates with 96 wells at a density of  $4 \times 10^4$  cells/well (100  $\mu$ l) in M199 medium containing 10% of bovine foetal serum and supplements (complete medium).

After incubation for 24 h (37°C, 5% CO<sub>2</sub>, 90% humidity) the cells were washed once and the medium was replaced with 200  $\mu$ l of complete medium containing the substances to be tested at final concentrations of  $10^{-5}$ ,  $10^{-6}$  and  $10^{-7}$  M. The test was performed in triplicate.

The plates were incubated for a further 48 h, after which 100  $\mu$ l of medium was removed and 20  $\mu$ l per well of dye solution was added in accordance with the supplier's instructions. The optical density ( $\lambda$  = 490 nm) was read after incubation for 1 h at 37°C with the use of a plate reader (Victor 2 - Wallace Perkin Elmer).

The results are expressed as the percentage inhibition of the formation of formazane in comparison with the control. The values obtained at the concentration of  $10^{-6}$  M are given in the following table.

Table 4: percentages of cytotoxicity towards HEP-G2 cells

Example	Cytotoxicity at $10^{-6}$ M
Ex. 01	3,8 %
Ex. 03	7,3 %
Ex. 04	0 %
Ex. 05	0 %
Ex. 06	13 %
Ex. 15	0 %

Example 24

Inhibition of the enzymatic activity of histone deacetylase

The capacity of the compounds of the invention to inhibit the activity of the histone deacetylase enzyme was evaluated with the use of the mouse HADC-1 enzyme. The assay was performed by the method already described in the literature [Biochem. Biophys. Acta, 1996, vol. 1296, p. 181]. The  $IC_{50}$  values (concentration which inhibits the activity of the enzyme by 50%) were derived from the percentages of inhibition obtained with various concentrations of the compound under test; these values are given in the following table.


Table 5: inhibition of mouse HDAC-1 enzyme

Example	IC <sub>50</sub> (nM)		Example	IC <sub>50</sub> (nM)
Ex. 01	142.0		Ex. 11	102.0
Ex. 03	219.0		Ex. 12	7.0
Ex. 04	113.0		Ex. 13	202.0
Ex. 05	90.0		Ex. 14	106.0
Ex. 06	19.0		Ex. 15	146.0
Ex. 07	14.0		Ex. 16	118.0
Ex. 08	58.0		Ex. 17	255.0
Ex. 09	67.0		Ex. 18	112.0
Ex. 10	130.0		Ex. 19	214.0



CLAIMS

1. Compounds of formula (I):



(I)

in which:

R is hydrogen, C<sub>1-4</sub> alkyl or phenyl;

R' is hydrogen or C<sub>1-4</sub> alkyl;

A is aryl or a monocyclic, dicyclic, or tricyclic residue, optionally partly or completely unsaturated, containing one or more heteroatoms selected from the group formed by N, S and O and possibly substituted with haloalkyl, alkylsulphonyl, (cycloalkyl)alkyl, alkanoyl, carbonyloxy, alkoxy, thioalkoxy, thiophenoxy, nitro, cyano, oxo, perfluoroalkoxy, perfluoroalkyl, phenyl, phenoxy, phenylalkoxy, benzoyloxy, phenylalkyl, benzoyl, phenylsulphonyl groups, the phenyl, phenoxy, phenylalkoxy, phenylalkyl, benzoyl and benzoyloxy substituents optionally being substituted in the aromatic ring with alkyl, alkoxy, alkylsulphonyl, amino, cyano, hydroxy, nitro, perfluoroalkoxy, perfluoroalkyl, phenylsulphonyl, thioalkoxy and halogen groups;

L is a chain of from 1 to 5 carbon atoms optionally containing a double bond or an NR' group in which R' is as defined above;

X is an oxygen atom, an NR' group, in which R' is as defined above, or is absent;

r and m are, independently, 0, 1 or 2;

B is phenyl or cyclohexyl.

2. Compounds according to Claim 1 in which R is hydrogen.

3. Compounds according to Claim 1 in which R is methyl.
4. Compounds according to any one of Claims 1-3 in which A is phenyl.
5. Compounds according to Claim 3 in which A is phenyl substituted with one or more groups selected from: alkoxy, nitro, perfluoroalkyl, phenoxy, phenyl, phenylalkoxy, benzyloxy, and thioalkoxy.
6. Compounds according to any one of Claims 1-5 in which X is absent.
7. Compounds according to any one of Claims 1-6 in which L is methylene or ethylene.
8. Compounds according to any one of Claims 1-7 in which m and r are equal to zero.
9. Compounds according to any one of Claims 1-8 in which B is phenyl.
10. Compounds according to any one of Claims 1-9 in which R' is hydrogen.
11. A compound selected from:
  - N-hydroxy-4-[2-(4-trifluoromethyl-phenyl)-acetylamino]-benzamide;
  - N-hydroxy-4-(3-phenyl-butyrylamino)-benzamide;
  - N-hydroxy-4-[3-(3-methoxy-phenyl)-propionylamino]-benzamide;
  - N-hydroxy-4-[2-(4-methoxy-phenyl)-acetylamino]-benzamide;
  - N-hydroxy-4-[2-(4-ethoxy-phenyl)-acetylamino]-benzamide;
  - N-hydroxy-4-[3-(3,5-bis-trifluoromethyl-phenyl)-propionylamino]-benzamide;
  - N-hydroxy-4-[2-(3,4,5-trimethoxy-phenyl)-acetylamino]-benzamide;
  - N-hydroxy-4-[2-(4-methylsulphanil-phenyl)-acetylamino]-benzamide;

N-hydroxy-4-[2-(3-trifluoromethyl-phenyl)-acetylamino]-benzamide;

N-hydroxy-4-[2-(3-nitro-phenyl)-acetylamino]-benzamide;

N-hydroxy-4-[2-(3-phenoxy-phenyl)-acetylamino]-benzamide;

N-hydroxy-4-[2-(diphenyl-4-yl)-acetylamino]-benzamide;

N-hydroxy-4-[2-(2,3-dimethoxy-phenyl)-acetylamino]-benzamide;

2-[2-(4-hydroxycarbamoyl-phenylcarbamoyl)-ethyl]-phenyl ester of benzoic acid;

N-hydroxy-4-[2-(4-nitro-phenyl)-acetylamino]-benzamide;

N-hydroxy-4-[2-(2-phenoxy-phenyl)-acetylamino]-benzamide;

N-hydroxy-4-[2-(4,5-dimethoxy-2-nitro-phenyl)-acetylamino]-benzamide;

N-hydroxy-4-[2-(2-benzyloxy-phenyl)-acetylamino]-benzamide;

N-hydroxy-4-[2-(2-nitro-phenyl)-acetylamino]-benzamide.

12. Compounds of any one of Claims 1-11 for use as medicaments.

13. Use of compounds of any one of Claims 1-11 for the preparation of medicaments with anti-inflammatory and/or auto-immune activity for the treatment of diseases such as spondyloarthropathy, rheumatoid arthritis, acute alcoholic hepatitis, inflammatory syndromes of the intestine (Crohn's disease and ulcerative colitis), asthma, diabetes, heart failure, intracerebral haemorrhage, psoriasis, atopic dermatitis, contact dermatitis, glomerulonephritis, systemic lupus erythematosus, chronic pulmonary obstruction, pulmonary fibrosis, multiple sclerosis, sepsis, septic shock, etc.

14. Use of the compounds of any one of Claims 1-11 for the preparation of medicaments for the treatment of tumorous and/or neurodegenerative diseases.
15. Use of the compounds of any one of Claims 1-11 and at least one active ingredient with anti-tumour action for the preparation of medicaments for the treatment of tumorous and neurodegenerative diseases.
16. Pharmaceutical compositions containing the compounds of any one of Claims 1-11 mixed with suitable excipients and/or vehicles.
17. Pharmaceutical compositions containing the compounds of any one of Claims 1-11 and at least one active ingredient with anti-tumour action mixed with suitable excipients and/or vehicles.

# INTERNATIONAL SEARCH REPORT

In International Application No.  
PCT/IT2004/000002

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07C259/10 A61K31/16 A61P35/00 A61P29/00

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07C A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, CHEM ABS Data

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 97/43251 A (BERTOLINI GIORGIO ;PAVICH GIANFRANCO (IT); BIFFI MAURO (IT); ITALF) 20 November 1997 (1997-11-20) cited in the application examples 1-10 page 30, line 2,3,12,13 page 31, line 29,30 page 32, line 4,5,10,11,16,17,22-31 page 33, line 1,2	1,2,4, 6-10,12, 13,16,17
Y	claims 4,5	1-17
X	WO 96/15105 A (ITALFARMACO SPA ;BERTOLINI GIORGIO (IT); AQUINO MARIO (IT); CHIAFF) 23 May 1996 (1996-05-23) examples 4,5	1,2, 7-10,12, 13,16,17
Y	claim 2	1-17
	--- -/-	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

3 May 2004

Date of mailing of the international search report

18/05/2004

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl  
Fax: (+31-70) 340-3016

Authorized officer

Fitz, W


# INTERNATIONAL SEARCH REPORT

Int'l Application No  
PCT/IT2004/000002

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>DATABASE CA 'Online! CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US; FUJII, SETSURO ET AL: "trans-4-Aminomethylcyclohexane hydroxamic acid" retrieved from STN Database accession no. 87:39013 XP002278214 abstract -&amp; JP 52 014747 A (ONO PHARMACEUTICAL CO., LTD., JAPAN) 3 February 1977 (1977-02-03)</p>	1,2,4,7, 10
Y	<p>UESATO S ET AL: "Novel histone deacetylase inhibitors: N-hydroxycarboxamides possessing a terminal bicyclic aryl group" BIOORGANIC &amp; MEDICINAL CHEMISTRY LETTERS, OXFORD, GB, vol. 12, 2002, pages 1347-1349, XP002276408 ISSN: 0960-894X page 1349; table 1 page 1347, column 1 page 1348, column 2, last line -page 1349, column 2</p>	1-17
A	<p>RICHON V M ET AL: "A CLASS OF HYBRID POLAR INDUCERS OF TRANSFORMED CELL DIFFERENTIATION INHIBITS HISTONE DEACETYLASES" PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF USA, NATIONAL ACADEMY OF SCIENCE. WASHINGTON, US, vol. 95, no. 6, 17 March 1998 (1998-03-17), pages 3003-3007, XP001120542 ISSN: 0027-8424 page 3003, column 1 page 3004; table 1</p>	1,11-17
A	<p>WO 02/055017 A (MISHRA NILAMADHAB ;KAMMER GARY M (US); UNIV WAKE FOREST (US)) 18 July 2002 (2002-07-18) claim 2</p>	1,11-17
P,X	<p>WO 03/013493 A (DINARELLO CHARLES ;BUFLER PHILIP (IT); DONA GIANCARLO (IT); ITALFA) 20 February 2003 (2003-02-20) page 12, Table, compounds 6,7,9-14 claim 1</p>	1,2,4, 6-10, 12-17

-/--

## INTERNATIONAL SEARCH REPORT

Int  I Application No  
PCT/IT2004/000002

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	BOUCHAIN, GILIANE ET AL: "Development of Potential Antitumor Agents. Synthesis and Biological Evaluation of a New Set of Sulfonamide Derivatives as Histone Deacetylase Inhibitors" JOURNAL OF MEDICINAL CHEMISTRY (2003), 46(5), 820-830 , XP002278212 page 823, table 3, compound 27 -----	1,2,7,9, 10,12, 14-17
P,X	LU, QIANG ET AL: "Zn <sup>2+</sup> -Chelating, Motif-Tethered, Short-Chain Fatty Acids as a Novel Class of Histone Deacetylase Inhibitors" JOURNAL OF MEDICINAL CHEMISTRY (2004), 47(2), 467-474 , 11 December 2003 (2003-12-11), XP002278213 page 471, Figure 3, cpds. -15,16,18,19,21,22 abstract -----	1,2,4, 6-10,12, 14-17

# INTERNATIONAL SEARCH REPORT

Information on patent family members

Int. Application No  
PCT/IT2004/000002

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9743251 A	20-11-1997	IT MI960968 A1	14-11-1997
		AU 713300 B2	25-11-1999
		AU 2896497 A	05-12-1997
		BR 9709234 A	10-08-1999
		CA 2254066 A1	20-11-1997
		CN 1221403 A ,B	30-06-1999
		CZ 9803667 A3	16-06-1999
		DE 69703207 D1	02-11-2000
		DE 69703207 T2	01-02-2001
		DK 901465 T3	18-12-2000
		WO 9743251 A1	20-11-1997
		EP 0901465 A1	17-03-1999
		ES 2151267 T3	16-12-2000
		GR 3035128 T3	30-04-2001
		JP 2000510472 T	15-08-2000
		KR 2000010982 A	25-02-2000
		PL 329873 A1	12-04-1999
		PT 901465 T	31-01-2001
		RU 2177473 C2	27-12-2001
		SK 157998 A3	13-04-1999
		US 6034096 A	07-03-2000
WO 9615105 A	23-05-1996	IT 1270694 B	07-05-1997
		AU 3927695 A	06-06-1996
		WO 9615105 A1	23-05-1996
		ZA 9509666 A	29-05-1996
JP 52014747 A	03-02-1977	NONE	
WO 02055017 A	18-07-2002	WO 02055017 A2	18-07-2002
		US 2003114525 A1	19-06-2003
		US 2003082666 A1	01-05-2003
WO 03013493 A	20-02-2003	IT MI20011733 A1	07-02-2003
		WO 03013493 A1	20-02-2003